

Stable systolic category of the product of spheres

HOIL RYU

The stable systolic category of a closed manifold M indicates the complexity in the sense of volume. This is a homotopy invariant, even though it is defined by some relations between homological volumes on M . We show an equality of the stable systolic category and the real cup-length for the product of arbitrary finite dimensional real homology spheres. Also we prove the invariance of the stable systolic category under the rational equivalences for orientable 0-universal manifolds.

57N65; 53C23, 55M30

1 Introduction

In this paper, a manifold is assumed to be closed, connected, orientable and smooth. The systole of a manifold M is the least length of non-contractible closed loops in M . One can generalize this concept to the least volume of k -dimensional nonzero homology classes, so called as the homology systole. Now we can imagine such systoles have some kind of relations with the entire volume of M , and it is natural to ask what kind of relationship exists.

As an answer, Gromov proved a theorem which says that the existence of non-trivial cup product implies the existence of the stable isosystolic inequality as follows.

Gromov's Theorem [6, 7.4.C] *Let M be an n -manifold (which can be non-orientable). If there exist some reduced real cohomology classes $\alpha_1^*, \dots, \alpha_k^*$ with α_i^* in $\tilde{H}^{d_i}(M; \mathbb{R})$ and a nonzero cup product $\alpha_1^* \cup \dots \cup \alpha_k^*$ in $\tilde{H}^n(M; \mathbb{R})$, then there exists $C > 0$ satisfying*

$$\prod_{i=1}^k \text{stsys}_{d_i}(M, \mathcal{G}) \leq C \cdot \text{mass}([M], \mathcal{G})$$

for all Riemannian metric $\mathcal{G}$ on M where stsys_{d_i} is the stable d_i -systole and $[M]$ is the fundamental class of M with coefficients in $\mathbb{Z}$ for orientable cases or $\mathbb{Z}/2\mathbb{Z}$ for non-orientable cases.

The greatest k satisfying the stable isosystolic inequality is called as the stable systolic category of M which is introduced by Katz and Rudyak [7], and it is known as a homotopy invariant by Katz and Rudyak [8]. We will show the stable systolic category of 0-universal manifold is also invariant under the rational equivalences in [Corollary 4.3](#).

For an orientable manifold M , Gromov's Theorem implies that the stable systolic category is not smaller than the real cup-length. So, is there some manifold M such that the stable systolic category is greater than the real cup-length? If such M exists, then the inversion of Gromov's Theorem will fail for M , while this interesting question is not answered yet. Instead of the answer, it is known the equality of them for some manifolds, eg, Dranishnikov and Rudyak [2]. In this paper, we also show more equality later in [Theorem 3.6](#) and [Theorem 3.8](#).

1.1 Definition of the stable systolic category

Let (X, A) be an object of the local Lipschitz category $\mathfrak{L}$. In this paper, we suppose X is a nonempty subset of some finite dimensional Euclidean space $\mathbb{R}^n$ with the standard norm. So X possesses the restricted metric of $\mathbb{R}^n$. Let G be a $\mathbb{Z}$ -module with a norm $|\cdot|$ which makes G a complete metric space. If G is $\mathbb{Z}$ or $\mathbb{R}$, we assume that norm of G is the standard norm. The comass of a differential form ω on X is defined as

$$\text{comass}(\omega) := \sup\{|\omega_x(\tau)| : x \in X, \text{ orthonormal } q\text{-frame } \tau\}.$$

Also, the mass of a q -current T in X is the dual norm of comass, ie,

$$\text{mass}(T) := \sup\{T(\omega) : \text{differential } q\text{-form } \omega, \text{comass}(\omega) \leq 1\}.$$

A Lipschitzian singular q -cube $\kappa : I^q \rightarrow X$, induces a homomorphism $\kappa_\flat$ from the module of polyhedral chains $\mathcal{P}_q(X; G)$ to the module of rectifiable currents $\mathcal{R}_q(X; G)$. Then the mass of κ is defined by the mass of the image $\kappa_\flat I^q$ where I^q is the corresponding polyhedral q -current of the unit rectangular parallelepiped I^q . This correspondence of κ to $\kappa_\flat I^q$ gives a chain map Φ of degree 0 from the chain complex of all Lipschitzian singular cubes into the chain complex of flat chains $\mathcal{F}_*(E|X)$. Then we can verify that Φ induces an isomorphism Φ_* between the flat homology module $H_q^b(X, A; G)$ and the singular homology module $H_q(X, A; G)$ for all q . One can find more details of these definitions at Federer [3], Federer [4], Federer and Fleming [5], Serre [9] and White [10].

For a Lipschitzian singular chain c , there exists a representation $\sum_i \kappa_i \otimes g_i$ where g_i is contained in G and κ_i is a Lipschitzian singular q -cube which is not overlapping

each other (subdivide if necessary). Then the *mass* of c is defined as

$$\text{mass}(c) := \sum_i |g_i| \cdot \text{mass}(\kappa_i).$$

The *mass* or *volume* of a singular homology class η in $H_q(X, A; G)$ is defined by

$$\text{mass}(\eta; G) := \inf \{ \text{mass}(c) : \eta = [c], c \text{ is a Lipschitzian cycle} \}.$$

If G is $\mathbb{R}$, the mass is a norm on the homology vector spaces. We will omit G in the case of $\mathbb{Z}$.

The q -dimensional homology systole of (X, A) is defined by infimum of mass of non-trivial q -th integral homology classes. However Gromov [6] claims that Gromov's Theorem will fail for $S^1 \times S^3$, if we consider the homology systoles instead of the stable systoles. Briefly, we can consider the stable systole as a systole in the real homology vector spaces. Here we give formal definition for the stable systole. The *stable mass* on $H_q(X, A; \mathbb{Z})$ is defined as

$$\text{stmass}(\eta) := \inf \{ \text{mass}(m \cdot \eta) / m : 0 < m \in \mathbb{Z} \}$$

for all $\eta \in H_q(X, A; \mathbb{Z})$. The inclusion $\iota : \mathbb{Z} \rightarrow \mathbb{R}$ induces the coefficient homomorphism ι_* on homology. Federer [4, 5.8] showed the stable mass of η is equal to the mass of the image $\iota_* \eta$. So we can define the q -dimensional stable systole of (X, A) as

$$\text{stsys}_q(X, A) := \inf \{ \text{stmass}(\eta) : \eta \in H_q(X, A; \mathbb{Z}), \iota_* \eta \neq 0 \}.$$

A homology q -systole or a stable q -systole is called *trivial*, if it is infinite. If the q -th real homology vector space $H_q(X, A; \mathbb{R})$ is zero, then the stable q -systole is trivial for all Riemannian metrics on (X, A) . Hence if the q -th integral homology module $H_q(X, A; \mathbb{Z})$ is a torsion module, then the stable q -systole is trivial for every metric on (X, A) .

For a given positive integer $n > 0$, a k -tuple $P = (p_1, \dots, p_k)$ of positive integers is called a *partition* of n if $n = p_1 + \dots + p_k$ and $p_1 \leq \dots \leq p_k \leq n$. A partition P is called *positive* (or *non-negative*) if $p_i > 0$ (or $p_i \geq 0$) for all i . The *size* of a partition which denoted by $\text{size}(P)$ is defined by the cardinality of positive integers contained in the partition. Hence if a k -tuple P is a positive partition, then the size of partition is k . From now on, we suppose a partition is positive unless otherwise stated. For a partition P , the *duplicated number* of p_i is the cardinality of elements in P who are equal to p_i .

Now we define concepts for an n -manifold M . A partition P of n is called *stable systolic categorical* for M , if there exists a real number $C > 0$ and non-trivial stable

p_i -systoles such that

$$\prod_{i=1}^{\text{size}(P)} \text{stsys}_{p_i}(M, \mathcal{G}) \leq C \cdot \text{mass}([M], \mathcal{G}; \mathbb{Z}/2\mathbb{Z})$$

for every Riemannian metric $\mathcal{G}$ on M where the fundamental class $[M]$ in $H_n(M; \mathbb{Z}/2\mathbb{Z})$.

Definition 1.1 The *stable systolic category* of M is defined by

$$\text{cat}_{\text{stsys}}(M) := \sup\{\text{size}(P) : P \text{ is stable systolic categorical partition for } M\}.$$

The *real cup-length* of M is defined by

$$\text{cup}_{\mathbb{R}}(M) := \min\{k \geq 0 : \alpha_0 \cup \alpha_1 \cup \cdots \cup \alpha_k = 0 \text{ for all } \alpha_i \in \tilde{H}^*(M; \mathbb{R})\}$$

where $\tilde{H}^*(M; \mathbb{R})$ is the reduced real cohomology ring of M . As we said before, the real cup-length is a lower estimate for the stable systolic category from Gromov's Theorem.

If M is non-orientable, then the top dimensional real cohomology vector space $H^n(M; \mathbb{R})$ vanishes. So every cohomology class in $H^n(M; \mathbb{R})$ vanishes, we can not apply Gromov's Theorem for top dimension. This is a reason to consider only orientable manifolds in this paper.

1.2 Acknowledgments

The author expresses gratitude to Professor Norio Iwase, whose guidance and support to write this paper.

2 Preliminaries on the stable systoles

Many equations and inequalities for mass are studied. One can find those results at Babenko [1], Federer [3] and Whitney [11]. Here we state or recall some of them for the stable systoles, with some appropriate modifications applied.

Proposition 2.1 For a local Lipschitz neighborhood retract X in $\mathbb{R}^n$, the stable 0-systole is 1.

Proof Let $\mathcal{D}_0(X)$ be the vector space of 0-currents. A map $\mathfrak{d}: X \rightarrow \mathcal{D}_0(X)$ can be defined as $\mathfrak{d}(x)(\omega) = \mathfrak{d}_x(\omega) := \omega(x)$ for a point x of X and a differential 0-form ω on X . Then $\mathfrak{d}_x$ is a polyhedral 0-current with $\text{mass}(\mathfrak{d}_x) = 1$. This implies that $\mathfrak{d}_x$ is a normal 0-cycle with coefficients $\mathbb{Z}$. Furthermore, the image $\iota_*\Phi_*^{-1}[\mathfrak{d}_x]$ is not vanished in $H_0(X; \mathbb{R})$. So we have

$$\text{stsys}_0(X) = \text{mass}(\iota_*\Phi_*^{-1}[\mathfrak{d}_x]) = 1$$

for an arbitrary point x in X . $\square$

Lemma 2.2 *For a local Lipschitz neighborhood retract X in $\mathbb{R}^n$, if one rescale the standard metric $\mathcal{G}$ on $\mathbb{R}^n$ by the square of a real number $t > 0$, then the quotient mass of a homology class $\eta \in H_q(X; G)$ increase by the t^q times. Furthermore, the stable q -systole satisfies*

$$\text{stsys}_q(X, t^2\mathcal{G}|X) = t^q \cdot \text{stsys}_q(X, \mathcal{G}|X)$$

where $\mathcal{G}|X$ is the restriction of $\mathcal{G}$ on X .

Proof A similar result was introduced by Whitney [11] for the real flat chains. So the first result is satisfied for an arbitrary homology class. Also the definition of the stable systole implies

$$\text{stsys}_q(X, t^2\mathcal{G}|X) = \inf \{t^q \cdot \text{mass}(\iota_*\eta, \mathcal{G}|X; \mathbb{R}) : \eta \in H_q(X, A; \mathbb{Z}), \iota_*\eta \neq 0\}$$

which means the equality for the stable systoles. $\square$

Proposition 2.3 [11, X.6 and X.7] *Let X and Y be open subsets in $\mathbb{R}^m$ and $\mathbb{R}^n$ respectively. For a locally Lipschitzian map $f: X \rightarrow Y$ and an integral rectifiable q -current T whose support is contained in a compact subset K of X , there exists an inequality*

$$\text{mass}(f_*T) \leq \text{Lip}(f|K)^q \cdot \text{mass}(T)$$

where $\text{Lip}(f|K)$ is the lower bound of Lipschitz constants of the restriction $f|K$.

Proposition 2.4 *Let (X, A) and (Y, B) be local Lipschitz neighborhood retract pairs in $\mathbb{R}^m$ and $\mathbb{R}^n$ respectively. If $f: (X, A) \rightarrow (Y, B)$ is a locally Lipschitzian map, then for any homology class η of $H_q(X, A; G)$, there is a compact subset K of $\mathbb{R}^m$ which satisfies*

$$0 \leq \text{mass}(f_*\eta; G) \leq \text{Lip}(f|K)^q \cdot \text{mass}(\eta; G)$$

where $f_*: H_q(X, A; G) \rightarrow H_q(Y, B; G)$ is the induced homomorphism.

Proof Note that f induces a homomorphism $f_b : Z_q(X, A; G) \rightarrow Z_q(Y, B; G)$ on flat cycles as well as $f_b \mathcal{F}_q(\mathbb{R}^m | A; G) \subset \mathcal{F}_q(\mathbb{R}^n | B; G)$. For a given flat homology class $\Phi_* \eta$, let T be a representative normal q -cycle in $Z_q(X, A; G)$. The naturality of Φ_* implies $\Phi_* f_* \eta = f_* \Phi_* \eta = f_* [T] = [f_b T]$. Also the relation of cosets $[f_b T] = [f_b T + f_b \mathcal{F}_q(\mathbb{R}^m | A; G)] = [f_b T + \mathcal{F}_q(\mathbb{R}^n | B; G)]$ implies that the relation of the sets

$$\{f_b T : [T] = \Phi_* \eta\} \subset \{S : [S] = \Phi_* f_* \eta\} \subset Z_q(Y, B; G).$$

With the definition of the mass of homology class, we obtain

$$\text{mass}(f_* \eta; G) \leq \inf \{\text{mass}(f_b T) : [T] = \Phi_* \eta\}.$$

Because of T is compact supported, there is a compact subset K of $\mathbb{R}^m$ with $\text{supp}(T) \subset \text{Int}(K)$. Here we can apply [Proposition 2.3](#) for T , so we have

$$\text{mass}(f_* \eta; G) \leq \text{Lip}(f|K)^q \cdot \inf \{\text{mass}(T) : [T] = \Phi_* \eta\}$$

which implies the result. $\square$

Lemma 2.5 *Let (X, A) and (Y, B) are local Lipschitz neighborhood retract pairs. If a locally Lipschitzian map $f : (X, A) \rightarrow (Y, B)$ induces a monomorphism $f_* : H_q(X, A; \mathbb{R}) \rightarrow H_q(Y, B; \mathbb{R})$, then there is a compact subset K in the ambient space of X satisfying*

$$\text{stsys}_q(Y, B) \leq \text{Lip}(f|K)^q \cdot \text{stsys}_q(X, A).$$

Furthermore, if $H_q(X, A; \mathbb{R})$ is nonzero, then $\text{stsys}_q(Y, B)$ is a positive real number.

Proof [Proposition 2.4](#) and $f_*(H_q(X, A; \mathbb{R}) \setminus \{0\}) \subset (H_q(Y, B; \mathbb{R}) \setminus \{0\})$ imply the existence of inequality in the stable systole level.

For integral homology class η with $\iota_* \eta$ is nonzero, the image $f_* \iota_* \eta$ does not vanish, since f_* is a monomorphism. Recall that the mass of real homology classes is a norm, hence $\text{mass}(f_* \iota_* \eta)$ is a positive real number. Furthermore, the stable q -systole does not converges to zero, since $\mathbb{Z}$ is discrete. $\square$

Proposition 2.6 *Let X and Y be open subsets of $\mathbb{R}^m$ and $\mathbb{R}^n$ respectively. For rectifiable currents S in $\mathcal{R}_p(X)$ and T in $\mathcal{R}_q(Y)$, the mass of their cross product is equal to the multiplication of their masses, ie,*

$$\text{mass}(S \times T) = \text{mass}(S) \cdot \text{mass}(T)$$

with respect to the product metric on $X \times Y$.

Proof Since S and T are rectifiable currents, mass can be written by associated Radon measures $\|S\|$, $\|T\|$ and $\|S \times T\|$. Therefore Fubini's Theorem (see Federer [3, 2.6.2.(2)]) implies

$$\text{mass}(S \times T) = \|S \times T\|(X \times Y) = \|S\|(X) \cdot \|T\|(Y) = \text{mass}(S) \cdot \text{mass}(T)$$

the result. $\square$

Lemma 2.7 *Let (X, A) and (Y, B) are local Lipschitz neighborhood retract pairs. For homology classes $\xi \in H_p(X, A; G)$ and $\eta \in H_q(Y, B; G)$, we can estimate*

$$\text{mass}(\xi \times \eta; G) \leq \text{mass}(\xi; G) \cdot \text{mass}(\eta; G)$$

$$\text{and} \quad \text{stsys}_{p+q}((X, A) \times (Y, B)) \leq \text{stsys}_p(X, A) \cdot \text{stsys}_q(Y, B)$$

with respect to the product metric on $(X, A) \times (Y, B)$.

Proof Let S and T be representative rectifiable cycles corresponding to ξ and η respectively, ie, $\Phi_*\xi = [S]$ with $S \in Z_p^b(X, A; G)$ and $\Phi_*\eta = [T]$ with $T \in Z_q^b(Y, B; G)$. Then the naturality of a cross product implies that there is a representative rectifiable current with the form of a cross product $S \times T$ in the coset $[c] = \Phi_*(\xi \times \eta)$. Therefore

$$\begin{aligned} \{S \times T : [S] \times [T] = \Phi_*\xi \times \Phi_*\eta\} &= \{S \times T : [S \times T] = \Phi_*(\xi \times \eta)\} \\ &\subset \{c : [c] = \Phi_*(\xi \times \eta)\} \\ &\subset Z_{p+q}^b((X, A) \times (Y, B); G). \end{aligned}$$

Hence [Proposition 2.6](#) implies an inequality

$$\begin{aligned} \text{mass}(\xi \times \eta; G) &\leq \inf\{\text{mass}(S \times T) : [S] \times [T] = \Phi_*\xi \times \Phi_*\eta\} \\ &= \text{mass}(\xi; G) \cdot \text{mass}(\eta; G) \end{aligned}$$

on homology level. To show the inequality of the stable systoles, recall that the cross product homomorphism

$$H_p(X, A; \mathbb{R}) \otimes H_q(Y, B; \mathbb{R}) \rightarrow H_{p+q}((X, A) \times (Y, B); \mathbb{R})$$

is a monomorphism. Therefore we can estimate the stable q -systole as

$$\begin{aligned} \text{stsys}_{p+q}((X, A) \times (Y, B)) &\leq \inf \left\{ \text{mass}(\xi \times \eta) : \begin{array}{l} \xi \in H_p(X, A; \mathbb{Z}), \iota_*\xi \neq 0, \\ \eta \in H_q(Y, B; \mathbb{Z}), \iota_*\eta \neq 0 \end{array} \right\} \\ &\leq \text{stsys}_p(X, A) \cdot \text{stsys}_q(Y, B). \end{aligned}$$

where the second inequality is obtained by the result on homology level. $\square$

Lemma 2.8 Suppose X and Y are local Lipschitz neighborhood retracts. If Y is connected and the Künneth formula gives an isomorphism of non-trivial vector spaces

$$H_q(X; \mathbb{R}) \otimes H_0(Y; \mathbb{R}) \cong H_q(X \times Y; \mathbb{R}) \neq \{0\},$$

then the stable q -systole satisfies

$$0 < \text{stsys}_q(X \times Y) = \text{stsys}_q(X) < \infty.$$

with respect to the product metric on $X \times Y$.

Proof Let $\text{pr}_1: X \times Y \rightarrow X$ be the first projection. From the assumption, for a nonzero homology class η in $H_q(X \times Y; \mathbb{R})$, there exist $[S] \neq 0$ in $H_q^b(X; \mathbb{R})$ and $[T] \neq 0$ in $H_0^b(Y; \mathbb{R})$ whose cross product is the image of η in $H_q^b(X \times Y; \mathbb{R})$ with the same positive mass, ie,

$$\text{mass}([S] \times [T]) = \text{mass}(\eta) > 0.$$

Note that the vector space of normal 0-chains $\mathcal{N}_0(Y; \mathbb{R})$ is equal to the vector space of polyhedral 0-chains $\mathcal{P}_0(Y; \mathbb{R})$ which is generated by $\{\partial_y : y \in Y\}$ where ∂ is defined in the proof of [Proposition 2.1](#). For every points y and y' in Y , $[\partial_y] = [\partial_{y'}]$ implies that there is a nonzero real number r such that $[T] = r[\partial_y]$ with $\text{mass}[T] = |r| \cdot \partial_y(1_Y^*) = |r|$. Also, every $[S] \times [T]$ has representation of $[r \cdot S] \times [\partial_y]$, therefore pr_{1*} is an isomorphism with $\text{pr}_{1*}([S] \times [T]) = [r \cdot S]$. Hence [Lemma 2.5](#) implies

$$\text{stsys}_q(X \times Y) \geq \text{stsys}_q(X) > 0$$

with the fact of pr_1 is a Lipschitzian map with $\text{Lip}(\text{pr}_1) = 1$. As a result, we obtain the equality by combining the result of [Lemma 2.7](#). $\square$

3 Calculation by dimension and constructing metrics

At first, we will calculate the stable systolic category from the dimensional information of homology. If the homology group is not so complex such as a real homology sphere, we know the stable systolic category by only using dimensional information. If a oriented manifold has a relatively simple cup-product structure such as n -fold producted space of spheres, then the stable systolic category can be also calculated instantly. Such methods to calculate the stable systolic category can be generalized as follows.

For a topological space X , let $\text{lpd}(X)$ denote the *least positive dimension* of real cohomology vector spaces of X . So $\text{lpd}(X) = l$ if and only if $\tilde{H}^i(X; \mathbb{R}) = \{0\}$ for $0 < i < l$ and $\tilde{H}^l(X; \mathbb{R}) \neq \{0\}$. If M is an m -manifold, then $\text{lpd}(M)$ is smaller than m .

Definition 3.1 An n -dimensional CW space X is said to have *maximal real cup length*, if there exist some real cohomology classes $\alpha_1, \dots, \alpha_r$ with $\alpha_i \in \tilde{H}^{d_i}(X; \mathbb{R})$, a nonzero cup-product $\alpha_1 \cup \dots \cup \alpha_r \in \tilde{H}^n(X; \mathbb{R})$ and $r := \lfloor n/\text{lpd}(X) \rfloor$ where $\lfloor x \rfloor$ denotes the floor of a real number x .

Example Let S be a manifold which is a real homology sphere. Then S has maximal real cup length, because of $\text{lpd}(S) = \dim(S)$. The n -fold direct product of S also has maximal real cup length. The direct product $S^2 \times S^3$ of spheres has maximal real cup length.

Corollary 3.2 If an m -manifold M has maximal real cup length, then the stable systolic category of M is equal to the real cup-length of M , ie,

$$\text{cat}_{\text{stsys}}(M) = \text{cup}_{\mathbb{R}}(M) = \lfloor m/\text{lpd}(M) \rfloor.$$

Proof We need to verify that $\text{cat}_{\text{stsys}}(M) \leq \text{cup}_{\mathbb{R}}(M)$. Let $r := \lfloor m/\text{lpd}(M) \rfloor$. If $(d_1, \dots, d_k)$ is a partition of m such that each stable d_i -systole is non-trivial, then $d_i \geq \text{lpd}(M)$, so there is an inequality

$$k \cdot \text{lpd}(M) \leq m = d_1 + \dots + d_k < (r+1) \cdot \text{lpd}(M)$$

which implies $k \leq r = \text{cup}_{\mathbb{R}}(M)$. $\square$

In general, the direct product $M \times N$ of manifolds does not have maximal real cup length even if M and N have maximal real cup-length. For example, the direct product of spheres $S^1 \times S^2$ does not have maximal real cup length.

Lemma 3.3 If manifolds $M_1^{m_1}, \dots, M_n^{m_n}$ have maximal real cup length, then the stable systolic category of their n -fold direct product $M_1 \times \dots \times M_n$ is greater than the sum of stable systolic categories for each M_i , ie,

$$\text{cat}_{\text{stsys}}(M_1 \times \dots \times M_n) \geq \text{cat}_{\text{stsys}}(M_1) + \dots + \text{cat}_{\text{stsys}}(M_n).$$

Proof Since M_i has maximal real cup length, there is nonzero cup product $\bigcup_{j=1}^{r_i} \alpha_i^j$ in $H^{m_i}(M_i; \mathbb{R})$ where $r_i := \lfloor m_i/\text{lpd}(M_i) \rfloor = \text{cat}_{\text{stsys}}(M_i)$ for $1 \leq i \leq n$.

By the Künneth formula, the n -fold cross product on the top dimensions induces an isomorphism

$$\bigotimes_{i=1}^n H^{m_i}(M_i; \mathbb{R}) \cong H^m \left(\prod_{i=1}^n M_i; \mathbb{R} \right) \quad \text{where} \quad m := \sum_{i=1}^n m_i.$$

This implies the existence of a nonzero cup product

$$\prod_{i=1}^n \left(\bigcup_{j=1}^{r_i} \alpha_i^j \right) = \bigcup_{i=1}^n \mathfrak{pr}_i^* \left(\bigcup_{j=1}^{r_i} \alpha_i^j \right) = \bigcup_{j=1}^{r_1} \mathfrak{pr}_1^* \alpha_1^j \cup \cdots \cup \bigcup_{j=1}^{r_n} \mathfrak{pr}_n^* \alpha_n^j$$

in the top-dimensional real cohomology vector space $H^m \left(\prod_{i=1}^n M_i; \mathbb{R} \right)$, where $\mathfrak{pr}_i : M_1 \times \cdots \times M_n \rightarrow M_i$ is the i -th projection. This cup product implies that $\text{cat}_{\text{sys}} \left(\prod_{i=1}^n M_i \right) \geq r_1 + \cdots + r_n$ by Gromov's Theorem. $\square$

Proposition 3.4 *For manifolds M and N , the least positive dimension of cohomology of $M \times N$ is the minimum of $\text{lpd}(M)$ and $\text{lpd}(N)$.*

Proof From the Künneth formula, $H^i(M \times N; \mathbb{R}) = \{0\}$ for $0 < i < \min(\text{lpd}(M), \text{lpd}(N))$. If $l := \min(\text{lpd}(M), \text{lpd}(N)) = \text{lpd}(M)$, then $H^l(M; \mathbb{R})$ is nonzero and the cross product homomorphism $H^l(M; \mathbb{R}) \otimes H^0(N; \mathbb{R}) \rightarrow H^l(M \times N; \mathbb{R})$ is a monomorphism. Therefore $H^l(M \times N; \mathbb{R})$ is nonzero. The case of $\text{lpd}(M) > \text{lpd}(N)$ is shown by using the same arguments. $\square$

For integers i and $j \neq 0$, let $\text{mod}(i, j)$ denotes the remainder from the division of i by j .

Corollary 3.5 *Suppose manifolds M^m and N^n have maximal real cup length, and an integer $l := \text{lpd}(M \times N)$. If M and N satisfy the conditions*

$$\lfloor m / \text{lpd}(M) \rfloor = \lfloor m / l \rfloor, \quad \lfloor n / \text{lpd}(N) \rfloor = \lfloor n / l \rfloor$$

and $\text{mod}(m, l) + \text{mod}(n, l) < l$,

then $M \times N$ has maximal real cup length. Therefore,

$$\text{cat}_{\text{sys}}(M \times N) = \text{cat}_{\text{sys}}(M) + \text{cat}_{\text{sys}}(N).$$

Proof Let integers $r := \lfloor m / l \rfloor$ and $s := \lfloor n / l \rfloor$.

[Proposition 3.4](#) implies that $l = \min(\text{lpd}(M), \text{lpd}(N)) = \text{lpd}(M \times N)$. So we can formulate $\lfloor (m+n) / \text{lpd}(M \times N) \rfloor = r + s + \lfloor \text{mod}(m, l) + \text{mod}(n, l) \rfloor$. By the assumption, $\lfloor \text{mod}(m, \text{lpd}(M)) + \text{mod}(n, \text{lpd}(N)) \rfloor$ is zero, so we have

$$\lfloor (m+n) / \text{lpd}(M \times N) \rfloor = r + s.$$

Thus it is sufficient to show that there is a nonzero cup product with the length of $r + s$.

Since M and N have maximal real cup length, there are cohomology classes $\alpha_1, \dots, \alpha_r$ and $\beta_1, \dots, \beta_s$ with their cup products are nonzero cohomology classes $\bigcup_{i=1}^r \alpha_i$ in $H^m(M; \mathbb{R})$ and $\bigcup_{i=1}^s \beta_i$ in $H^n(N; \mathbb{R})$. From the proof of [Lemma 3.3](#), there is a nonzero cup product $\bigcup_{i=1}^r \mathfrak{pr}_1^* \alpha_i \cup \bigcup_{i=1}^s \mathfrak{pr}_2^* \beta_i$ in the top dimensional cohomology vector space $H^{m+n}(M \times N; \mathbb{R})$. $\square$

Without the condition of the product $M \times N$ has maximal real cup length, we can generalize this corollary as follow.

Theorem 3.6 *Let manifolds M^m and N^n have maximal real cup length. If*

$$\text{mod}(m, \text{lpd}(M)) + \text{mod}(n, \text{lpd}(N)) < \max(\text{lpd}(M), \text{lpd}(N)),$$

then the stable systolic category of their product $M \times N$ is the sum of each stable systolic category, ie,

$$\text{cat}_{\text{stsys}}(M \times N) = \text{cat}_{\text{stsys}}(M) + \text{cat}_{\text{stsys}}(N).$$

Proof Since M and N have maximal real cup length,

$$r := \lfloor m / \text{lpd}(M) \rfloor = \text{cat}_{\text{stsys}}(M) \quad \text{and} \quad s := \lfloor n / \text{lpd}(N) \rfloor = \text{cat}_{\text{stsys}}(N).$$

In the case of $\text{lpd}(M) = \text{lpd}(N)$ is [Corollary 3.5](#). So we will assume $\text{lpd}(M) < \text{lpd}(N)$.

From [Lemma 3.3](#), $\text{cat}_{\text{stsys}}(M \times N) \geq \text{cat}_{\text{stsys}}(M) + \text{cat}_{\text{stsys}}(N) = r + s$. Therefore, it is sufficient to show that any partition of $m+n$ whose size is greater than $r+s$, is not a stable systolic categorical partition.

Suppose $m+n = d_1 + \dots + d_k$ is a stable systolic categorical partition for $M \times N$ with some integer $1 \leq r' \leq k$ and the condition $0 < \text{lpd}(M) \leq d_1 \leq \dots \leq d_{r'} < \text{lpd}(N)$. For an arbitrary $t \geq 1$, let $\mathcal{G}_t := t^2 \mathcal{G}_M + \mathcal{G}_N$ be a Riemannian metric on $M \times N$. Then [Lemma 2.2](#) and [Lemma 2.8](#) imply that the stable systoles for the partition $d_1 + \dots + d_k$ satisfies

$$\begin{aligned} \prod_{i=1}^k \text{stsys}_{d_i}(M \times N, \mathcal{G}_t) &\geq \prod_{i=1}^{r'} \text{stsys}_{d_i}(M, t^2 \mathcal{G}_M) \prod_{j=r'+1}^k \text{stsys}_{d_j}(M \times N, \mathcal{G}_t) \\ &= t^{d_1 + \dots + d_{r'}} \prod_{i=1}^{r'} \text{stsys}_{d_i}(M, \mathcal{G}_M) \prod_{j=r'+1}^k \text{stsys}_{d_j}(M \times N, \mathcal{G}_t) \end{aligned}$$

Since we assume that $t \geq 1$, we can obtain the inequality $\text{stsys}_{d_j}(M \times N, \mathcal{G}_t) \geq \text{stsys}_{d_j}(M \times N, \mathcal{G}_1)$ for each $r' + 1 \leq j \leq k$. On the other hands, the mass of integral fundamental class $[M \times N]$ is characterized by [Lemma 2.2](#) and [Lemma 2.7](#) as

$$\begin{aligned} \text{mass}([M \times N], \mathcal{G}_t) &\leq \text{mass}([M], t^2 \mathcal{G}_M) \cdot \text{mass}([N], \mathcal{G}_N) \\ &= t^m \cdot \text{mass}([M], \mathcal{G}_M) \cdot \text{mass}([N], \mathcal{G}_N). \end{aligned}$$

Here if we assume that $d_1 + \dots + d_{r'} > m$, then we have

$$\frac{\prod_{i=1}^k \text{stsys}_{d_i}(M \times N, \mathcal{G}_t)}{\text{mass}([M \times N], \mathcal{G}_t)} \geq t^{m - (d_1 + \dots + d_{r'})} \cdot \frac{\prod_{i=1}^{r'} \text{stsys}_{d_i}(M, \mathcal{G}_M) \cdot \prod_{j=r'+1}^k \text{stsys}_{d_j}(M \times N, \mathcal{G}_1)}{\text{mass}([M], \mathcal{G}_M) \cdot \text{mass}([N], \mathcal{G}_N)}$$

where the right-hand side of the inequality diverges as $t \rightarrow \infty$. This contradicts to that $(d_1, \dots, d_k)$ is a stable systolic categorical partition. Hence we obtain $d_1 + \dots + d_{r'} \leq m$ and $d_{r'+1} + \dots + d_k \geq n$. This condition for m implies

$$r' \leq \lfloor (d_1 + \dots + d_{r'}) / \text{lpd}(M) \rfloor \leq \lfloor m / \text{lpd}(M) \rfloor \leq r.$$

Let $s' := k - r'$. From the assumption, $\text{lpd}(M) / \text{lpd}(N) < 1$ and

$$\text{mod}(m, \text{lpd}(M)) + \text{mod}(n, \text{lpd}(N)) < \text{lpd}(N),$$

so we can calculate as

$$k = r' + s' \leq r + s$$

which implies $\text{cat}_{\text{stsys}}(M \times N) \leq \text{cat}_{\text{stsys}}(M) + \text{cat}_{\text{stsys}}(N)$. $\square$

Corollary 3.7 Suppose manifolds $M_0 \times M_1 \times \dots \times M_k$ and $M_{k+1} \times \dots \times M_n \times M_{n+1}$ have maximal real cup length with

$$\text{lpd}(M_0) = \text{lpd}(M_1) = \dots = \text{lpd}(M_k)$$

$$\text{and} \quad \text{lpd}(M_{k+1}) = \dots = \text{lpd}(M_n) = \text{lpd}(M_{n+1}).$$

Let $r_i := \lfloor \dim(M_i) / \text{lpd}(M_i) \rfloor$ for $0 \leq i \leq n+1$. If $M_0, \dots, M_{n+1}$ satisfy conditions $\dim(M_i) = \text{lpd}(M_i) \cdot r_i$ for $1 \leq i \leq n$ and

$$\begin{aligned} \dim(M_0) - \text{lpd}(M_0) \cdot r_0 + \dim(M_{n+1}) - \text{lpd}(M_{n+1}) \cdot r_{n+1} \\ < \max(\text{lpd}(M_0), \text{lpd}(M_{n+1})), \end{aligned}$$

then:

$$\text{cat}_{\text{stsys}}\left(\prod_{i=0}^{n+1} M_i\right) = \sum_{i=0}^{n+1} \text{cat}_{\text{stsys}}(M_i) = \sum_{i=0}^{n+1} r_i$$

For the product $S^1 \times S^2$ of spheres, [Theorem 3.6](#) can not applied. So we must approach from the other viewpoints to obtain the stable systolic category.

Theorem 3.8 If manifolds $S_1^{m_1}, \dots, S_n^{m_n}$ are real homology spheres, then the stable systolic category of their n -fold direct product is the number of spheres.

Proof Since every real homology spheres have maximal real cup length, [Lemma 3.3](#) gives us a lower bound $\text{cat}_{\text{stsys}}(S_1 \times \dots \times S_n) \geq n$.

Suppose $m_i \leq m_{i+1}$ for each $1 \leq i \leq n$. Then a partition $(m_1, \dots, m_n)$ of $\sum_i m_i$ can be rewritten as $(r_1, \dots, r_1, r_2, \dots, r_{l-1}, r_l, \dots, r_l)$ where r_i is a range. This corresponding to rewrite

$$S_1^{m_1} \times \dots \times S_n^{m_n} = (S_1^{r_1} \times \dots \times S_{s_1}^{r_1}) \times (S_{s_1+1}^{r_2} \times \dots \times S_{s_1+s_2}^{r_2}) \times \dots \\ \times (S_{s_1+\dots+s_{l-1}+1}^{r_l} \times \dots \times S_{s_1+\dots+s_{l-1}+s_l}^{r_l})$$

where $r_i := m_{s_1+\dots+s_{i-1}+1} = \dots = m_{s_1+\dots+s_{i-1}+s_i}$ with $r_i < r_{i+1}$ and $s_i > 0$ is the duplicated number of r_i , so that $s_1 + \dots + s_l = n$. For simplicity, let

$$X_p := S_1 \times \dots \times S_{s_1+\dots+s_p} \quad \text{and} \quad Y_p := S_{s_1+\dots+s_p} \times \dots \times S_n$$

for $1 \leq p \leq n$. Then $S_1 \times \dots \times S_n = X_p \times Y_p$ and we can observe that $\mathcal{G}_{p,t} := t^2 \mathcal{G}_{X_p} + \mathcal{G}_{Y_p}$ is a Riemannian metric on $X_p \times Y_p$ for $t > 0$ when $\mathcal{G}_{X_p} + \mathcal{G}_{Y_p}$ is a Riemannian metric on $X_p \times Y_p$. Now we can apply [Lemma 2.8](#) and [Lemma 2.2](#), so

$$\text{stsys}_q(X_p \times Y_p, \mathcal{G}_{p,t}) = \text{stsys}_q(X_p, t^2 \mathcal{G}_{X_p}) = t^q \cdot \text{stsys}_q(X_p, \mathcal{G}_{X_p})$$

for the non-trivial stable systoles in the dimension of $1 \leq q \leq s_1 + \dots + s_p$.

Here we suppose $(d_1, \dots, d_k)$ with $d_i \leq d_{i+1}$, is the longest stable systolic categorical partition for $S_1 \times \dots \times S_n$. Then we can rewrite $(d_1, \dots, d_k)$ by the ranges $\{r_1, \dots, r_l\}$ and the duplicated number $s'_i \geq 0$ of r_i . We will show this partition is not longer than n by induction on p for $1 \leq p \leq l$ and contradiction. Assume that $s'_i = s_i$ for $1 \leq i \leq p-1$. If $s'_p > s_p$, then using a similar argument in the proof of [Theorem 3.6](#), we can observe that the right-hand side of the inequality

$$\frac{\prod_{i=1}^k \text{stsys}_{d_i}(X_p \times Y_p, \mathcal{G}_{p,t})}{\text{mass}([X_p \times Y_p], \mathcal{G}_{p,t})} \geq t^w \cdot \frac{\prod_{i=1}^p \text{stsys}_{r_i}(X_p, \mathcal{G}_{X_p})^{s'_i} \prod_{i=p+1}^l \text{stsys}_{r_i}(X_p \times Y_p, \mathcal{G}_{p,1})^{s'_i}}{\text{mass}([X_p], \mathcal{G}_{X_p}) \cdot \text{mass}([Y_p], \mathcal{G}_{Y_p})}$$

diverge as $t \rightarrow \infty$ where $w := r_1(s'_1 - s_1) + \dots + r_l(s'_l - s_l) > 0$. This contradicts to that the partition $(d_1, \dots, d_k)$ is stable systolic categorical, and hence we obtain $s'_p \leq s_p$. However we must choose $s'_p = s_p$ to make the longest partition. As a result, the size of the longest stable systolic categorical partition can not exceed $n = s_1 + \dots + s_l$. $\square$

4 Invariance under the rational equivalences

Suppose M and N are n -manifolds. Let K and L be a triangulation of M and N respectively. In this section, K and L are subdivided if necessary, but we will use the same symbol. For a continuous map $f: M \rightarrow N$, there is a non-degenerate simplicial

approximation $g: K \rightarrow L$ of f . For an open n -simplex e in L , consider a map $h: K \xrightarrow{g} L \rightarrow L/(L \setminus e)$. We will call $\deg(h)$ the *degree* of g at e which is denoted by $\deg_e(g)$. Let

$$D(g) := \sup\{|\deg_e(g)| : \text{open } n\text{-simplex } e \text{ in } L\}.$$

Here $D(g)$ is finite, because of we can assume that K and L are finite simplicial complexes.

For an arbitrary Riemannian metric $\mathcal{G}_N$ on N and $\varepsilon > 0$, there is a piecewise linear metric $\mathcal{G}_L = \mathcal{G}_L(\varepsilon)$ on L satisfying

$$|\text{stsys}_q(L, \mathcal{G}_L) - \text{stsys}_q(N, \mathcal{G}_N)| \leq \varepsilon$$

for every non-trivial stable q -systoles (compare Federer [3, 4.1.22]) and the realization of L with $\mathcal{G}_L$ is a PL section of the normal bundle over N with $\mathcal{G}_N$ in some finite dimensional Euclidean space. Such metric can be obtained by subdividing K and L , and translating vertices in L along the fiber of the normal bundle to do not degenerate any simplex. For $0 < \varepsilon' < \varepsilon$, a suitable metric $\mathcal{G}_L(\varepsilon')$ also can be acquired by the same way. Hence we can assume that $D(g)$ is not changed by ε and $\mathcal{G}_L$. As ε approaches to 0, each L , $\mathcal{G}_L$ and $g^*\mathcal{G}_L$ converges to N , $\mathcal{G}_N$ and a piecewise Riemannian metric on M respectively. Under this circumstance, we obtain following lemma.

Lemma 4.1 *Suppose q -th real homology vector space of K and L are non-trivial. If $g: K \rightarrow L$ induces a monomorphism g_* between the q -th real homology vector spaces, then*

$$\text{stsys}_q(L, \mathcal{G}_L) \leq \text{stsys}_q(K, g^*\mathcal{G}_L) \leq D(g) \cdot \text{stsys}_q(L, \mathcal{G}_L) < \infty$$

for every piecewise linear metric $\mathcal{G}_L$ on L .

Proof With the pullback PL metric $g^*\mathcal{G}_L$ on K , g is a distance decreasing map. Combining this with Lemma 2.5,

$$\text{stsys}_q(L, \mathcal{G}_L) \leq \text{Lip}(g)^q \cdot \text{stsys}_q(K, g^*\mathcal{G}_L) \leq \text{stsys}_q(K, g^*\mathcal{G}_L).$$

On the other hands, the inverse image of an arbitrary q -simplex of L is $D(g)$ of q -simplices at most, since g is a non-degenerate simplicial map and every q -simplex is contained in the boundary of some n -simplex for $q < n$. Also each simplex in the inverse image has same mass of the preimage, since the restriction of g on each simplex is isometry. This implies that the mass of a q -chain c of K is not greater than $D(g)$ times of the mass of the image $g_b(c)$ which is not trivial. Therefore we can verify that

$$\text{stsys}_q(K, g^*\mathcal{G}_L) \leq D(g) \cdot \text{stsys}_q(L, \mathcal{G}_L)$$

for an arbitrary PL metric $\mathcal{G}_L$. □

Remark If K is not a triangulation of a manifold, we can not sure that every q -simplex of K is contained in the boundary of some n -simplex for $q < n$. For example, a triangulation of the one-point union $S^1 \vee S^2$ has some 1-simplex in S^1 which is not contained in the boundary of any 2-simplex.

Since the stable systolic category is a homotopy invariant, here we obtain following proposition using similar techniques of Katz and Rudyak [8].

Proposition 4.2 *Let M and N are n -manifolds. If there exists a smooth map $f: M \rightarrow N$ which induces a monomorphism on every real homology vector space, then $\text{cat}_{\text{sys}}(M) \leq \text{cat}_{\text{sys}}(N)$.*

Proof We apply Lemma 4.1,

$$\text{sys}_q(N, \mathcal{G}_N) \leq \text{sys}_q(L, \mathcal{G}_L) + \varepsilon \leq \text{sys}_q(K, g^* \mathcal{G}_L) + \varepsilon$$

$$\text{and } \text{sys}_q(N, \mathcal{G}_N) + \varepsilon \geq \text{sys}_q(L, \mathcal{G}_L) \geq 1/D(g) \cdot \text{sys}_q(K, g^* \mathcal{G}_L)$$

where L converges to N in some Euclidean space and $g^* \mathcal{G}_L$ converges to a piecewise Riemannian metric $\mathcal{G}_M$ on M as ε approaches to 0. Suppose there exists a stable systolic categorical partition $(d_1, \dots, d_k)$ for M . Then there exist $C > 0$ and $\delta = \delta(\varepsilon) > 0$ such that δ converges to 0 as ε approaches to 0 and

$$\prod_{i=1}^k \text{sys}_{d_i}(K, g^* \mathcal{G}_L) \leq C \cdot \text{mass}([K], g^* \mathcal{G}_L) + \delta,$$

because of each metric $g^* \mathcal{G}_L$ can be approximated by some Riemannian metrics on M . We can assume that $\varepsilon \leq \text{sys}_{d_i}(N, \mathcal{G}_N)$ for all i , so

$$\begin{aligned} \prod_{i=1}^k \text{sys}_{d_i}(L, \mathcal{G}_L) &\leq 2^k \cdot \prod_{i=1}^k \text{sys}_{d_i}(K, g^* \mathcal{G}_L) \\ &\leq 2^k C \cdot \text{mass}([K], g^* \mathcal{G}_L) + 2^k \delta \\ &\leq 2^k CA(g) \cdot \text{mass}([L], \mathcal{G}_L) + 2^k (CA(g) \cdot \varepsilon + \delta). \end{aligned}$$

This implies the partition $(d_1, \dots, d_k)$ is also stable systolic categorical for N . Therefore we obtain the result $\text{cat}_{\text{sys}}(M) \leq \text{cat}_{\text{sys}}(N)$. $\square$

Let X and Y are simply connected spaces. A continuous map $f: X \rightarrow Y$ is called a *rational equivalence*, if the induced map $f^*: H^*(Y; \mathbb{Q}) \rightarrow H^*(X; \mathbb{Q})$ is an isomorphism.

Corollary 4.3 *The stable systolic category of a 0-universal manifold is invariant under the rational equivalences.*

Proof For a 0-universal manifold M and a rational equivalence to a space X , there exists a rational equivalence from X to M . $\square$

References

- [1] **I K Babenko**, *Asymptotic invariants of smooth manifolds*, Russian Acad. Sci. Izv. Math. 41 (1993) 1–38
- [2] **A N Dranishnikov, Y B Rudyak**, *Stable systolic category of manifolds and the cup-length*, J. Fixed Point Theory Appl. 6 (2009) 165–177
- [3] **H Federer**, *Geometric measure theory*, Grundlehren der mathematischen Wissenschaften 153, Springer (1969)
- [4] **H Federer**, *Real flat chains, cochain and variational problems*, Indiana Univ. Math. J. 24 (1974) 351–407
- [5] **H Federer, W H Fleming**, *Normal and integral currents*, Ann. Math. 72 (1960) 458–520
- [6] **M Gromov**, *Filling Riemannian manifolds*, J. Differential Geom. 18 (1983) 1–147
- [7] **M G Katz, Y B Rudyak**, *Lusternik-Schnirelmann category and systolic category of low dimensional manifolds*, Comm. Pure Appl. Math 59 (2006) 1433–1456
- [8] **M G Katz, Y B Rudyak**, *Bounding volume by systoles of 3-manifolds*, J. London Math. Soc. 78 (2008) 407–417
- [9] **J P Serre**, *Homologie singulière des espaces fibrés*, Ann. Math. 54 (1951) 425–505
- [10] **B White**, *Rectifiability of flat chains*, Ann. Math. 150 (1999) 165–184
- [11] **H Whitney**, *Geometric integration theory*, Annals of Mathematics Studies 21, Princeton University Press (1957)

Graduate School of Mathematics, Kyushu University
774, Motoooka, Nishi-ku, Fukuoka, 819-0395, Japan

h-ryu@math.kyushu-u.ac.jp